

Preliminary results from the AROMAT 2014 campaign



A. Merlaud & Aromat Team, DOAS seminars, 27 March 2015


Outline

- Context
- Presentation of the campaign
- Preliminary results
- Preliminary conclusions and perspectives

Collaboration with the University of Galati since 2011

Development and tests of the SWING-UAV payload (BIRA), and of a flying wing UAV (Uni. Galati and Reev River Aerospace)

Mobile-DOAS studies: Constantin et al., Sensors, 2013, Dragomir et al. Atmos Pol Res, 2015



Hotspots near Galati



Galati Arcelor Mittal Steel factory

Slobozia Fertilizer factory



Romania recently entered in ESA (Dec. 2011)

ROMANIAN FLAG RAISED AT ESA



24 January 2012 The New Year sees the Romanian flag now flying at ESA sites alongside those of ESA's other Member States, after Romania officially became ESA's 19th Member State on 22 December.

The Agreement on Romania's accession to the ESA Convention was signed on 20 January 2011 in Bucharest, by Jean-Jacques Dordain, Director General of ESA, Teodor Baconschi, Romanian Minister of Foreign Affairs, and Marius-Ioan Piso, President and CEO of the Romanian Space Agency.



contribution




georeturn



Sentinel 5p mission

TROPOMI scheduled to be launched in 2016, on an Eurockot, from Plesetsk Cosmodrome




Outline

- Context
- Presentation of the campaign
- Preliminary results
- Preliminary conclusions and perspectives

AROMAT: Airborne Romanian Measurements of Aerosols and Trace gases

- ESA campaign coordinated by BIRA-IASB (Belgium), with participation of 7 other research institutions: Uni. Galati, INOE and INCAS (Romania), IUP-Bremen, Uni. Berlin, MPIC (Germany), and KNMI (The Netherlands)
- Aim: test newly developed airborne instruments for air quality research and prepare forthcoming intercomparison/S5P validation campaign(s) in Romania
- First week of September 2014 in Bucharest (largest city in Romania, with 2.2 million pop., heavy traffic)
- Second week in Turceni: small village on the countryside, with a very large power plant known to be a major source of NO_x, SO₂ and aerosols

→ Both spots visible on OMI data →



Instruments and platforms

Airborne:

- Uni. Berlin Cessna:
→ Sunphotometer and Microtops (Uni. Berlin) 
→ AirMap (IUP-Bremen) 
- Weather balloons
→ NO₂ sonde (KNMI) 
- INCAS UAV
→ Dust trak (INCAS) 
→ Aerosol particle sizer (INOE) 
- Uni. Galati UAV
→ SWING (BIRA) 
→ NO₂ sonde (KNMI) 

Mobile (car-based)

- Mini-MaxDOAS (MPIC) 
- Zenith-only mobile DOAS (Uni. Galati, BIRA) 
- Double channel mobile DOAS (BIRA) 
- NO₂ sonde (KNMI) 

NO₂  SO₂  aerosols 

Static/Movable

- RADO observatory (Bucharest): lidars, in situ monitors, Aerosol Chemical Speciation Monitor, meteo. Station, PM 2.5, Aethalometer, CIMEL sun photometer (INOE) 
- In situ monitor (Uni. Galati) 
- In situ monitor, scanning lidar, ACSM (INOE) 

Airborne Platforms in AROMAT-2014



INCAS UAV



FUB Cessna



UGAL/RRA UAV



KNMI Weather balloon

Mobile Platforms in AROMAT-2014



Mobile DOAS (5 spectrometers),
MPIC, UGAL, BIRA




Mobile lab (INOE, UGAL)

Outline

- Context
- Presentation of the campaign
- Preliminary results
- Preliminary conclusions and perspectives


Bucharest



Bucharest: INOE APS on INCAS UAV





APS: Aerosol Particle Sizer




Bucharest: Mobile DOAS




Intercomparison with MPIC and UGAL (Sept 2)




Bucharest: Mobile DOAS Intercomparison with MPIC and UGAL (Sept 3)



Bucharest: Measurements at RADO observatory (INOE)



Bucharest: Measurements at RADO observatory (INOE)



Magurele, 01 Sep. 2014 – one hour average

Bucharest: Coincident AirMAP and Mobile DOAS measurements (Sept 8)





Fig 5a: Comparison between AirMAP and UGAL Car-DOAS NO_2 DSCDs:

Pixel-wise correlation plot of NO_2 differential slant column densities (DSCDs) of gridded datasets ($0.001^\circ \times 0.001^\circ \approx 100 \text{ m}^2$) of co-located measurements of AirMAP pixels and car positions.




Fig 5b: Time series of the car-DOAS measurements. Additionally shown are the corresponding AirMAP measurements for the for car locations. The plot at the bottom shows the time difference between the measurements.

*Andreas Meier, poster at
the German Physical Society*





Image © 2015 CNES / Astrium


© 2015 Google

Google earth


Turceni Power plant












Turceni: Lidar measurements






Turceni: NO₂ sonde measurements




Turceni: NO₂ sonde measurements



11 vertical
NO₂ profiles,
all flown from
Turceni



Mobile DOAS measurements around Turceni



10 SEPTEMBER 2014 - COMPARISON WITH OMI (13.09)




OMI DETECT $\sim 6 \times 10^{15}$ molec./cm²


While inside the pixel (13x24km²) mobile DOAS “see” 1×10^{17} molec/cm²



Mobile DOAS measurements around Turceni






Mobile DOAS measurements around Turceni, SO₂





3. Preliminary results of AROMAT flights

Flight 2- Sept 11






3. Preliminary results of AROMAT flights

Flight 2- Sept 11



3. Preliminary results of AROMAT flights





Outline

- Context
- Presentation of the campaign
- Preliminary results
- Preliminary conclusions and perspectives


Preliminary conclusions from AROMAT-1

- all the airborne platforms/instruments operated
- First successful MobileDOAS intercomparison for NO₂
- In-situ intercomparison more difficult
- AirMap seems most promising for satellite validation in the short term
- Mobile DOAS meas. are useful to plan the flights/validate the airborne meas.
- First SWING-UAV data with a clear signal and spatial gradients, but georeferencing is difficult
- Turceni not the easiest place for the sondes on balloon
- UAVs are dangerous
- Jiu Valley not optimal for satellite validation but very good to test new instruments

Perspectives

UAV flights above
the Black Sea for
monitoring ship
emissions (NO₂ and
SO₂, with a new
spectrometer)

-> June 2015



Perspectives

- AROMAT-2 in Aug/Sept 2015
 - > Addition (?) of a third airborne imager: APEX, operated from the DLR Dornier
 - > KNMI sondes onboard a manned aircraft to get profile around Bucharest
 - > GB Imaging from the ground in Turceni (SO₂ camera and Altius breadboard for NO₂)
- Bucharest campaign in summer 2017